

CORRECTION

Open Access



Correction to: Plasma fatty acid levels and gene expression related to lipid metabolism in peripheral blood mononuclear cells: a cross-sectional study in healthy subjects

Sunniva V. Larsen¹, Kirsten B. Holven^{1,2}, Inger Ottestad¹, Kine N. Dagsland³, Mari C. W. Myhrstad³ and Stine M. Ulven^{1*}

Correction

Unfortunately, after publication of this article [1], it was noticed that some corrections to Table 2 were not carried out. The corrected Table 2 can be seen below and the original article has also been updated to reflect this.

Author details

¹Department of Nutrition, Institute for Basic Medical Sciences, University of Oslo, P.O. Box 1046, Blindern, 0317 Oslo, Norway. ²Norwegian National Advisory Unit on Familial Hypercholesterolemia, Department of Endocrinology, Morbid Obesity and Preventive Medicine, Oslo University Hospital, P.O. Box 4950, Nydalen, 0424 Oslo, Norway. ³Department of Health, Nutrition and Management, Faculty of Health Sciences, Oslo and Akershus University College of Applied Sciences, P.O. Box 4, St. Olavs plass, 0130 Oslo, Norway.

Received: 20 June 2018 Accepted: 20 June 2018

Published online: 05 July 2018

Reference

1. Larsen SV, Holven KB, Ottestad I, Dagsland KN, Myhrstad MCW, Ulven SM. Plasma fatty acid levels and gene expression related to lipid metabolism in peripheral blood mononuclear cells: a cross-sectional study in healthy subjects. *Genes Nutr.* 2018;13:9. <https://doi.org/10.1186/s12263-018-0600-z>.

* Correspondence: smulven@medisin.uio.no

¹Department of Nutrition, Institute for Basic Medical Sciences, University of Oslo, P.O. Box 1046, Blindern, 0317 Oslo, Norway

Full list of author information is available at the end of the article



Table 2 Characteristics and plasma fatty acid profile of subjects in the highest and lowest tertiles

	n-6 level			n-3 level			SFA/PUFA ratio		
	Highest tertile	Lowest tertile	<i>P</i>	Highest tertile	Lowest tertile	<i>P</i>	Highest tertile	Lowest tertile	<i>P</i>
	(n 18)	(n 18)		(n 18)	(n 18)		(n 18)	(n 18)	
Male/female	5/13	5/13		3/15	5/13		5/13	6/12	
Age (years)	25 (23–33)	25 (21–29)	0.31	25 (21–31)	24 (21–28)	0.80	25 (22–29)	28 (23.0–32.8)	0.38
BMI (kg/m ²)	22.5 ± 2.9	22.6 ± 2.6	0.94	22.2 ± 2	22.7 ± 3.3	0.57	23.3 ± 2.7	22.6 ± 2.6	0.41
Total-C (mmol/l)	4.9 ± 0.9	4.7 ± 1.1	0.65	4.9 ± 1	4.9 ± 1	0.96	4.7 ± 0.8	5.0 ± 0.9	0.25
LDL-C (mmol/l)	2.7 ± 0.8	2.6 ± 0.9	0.79	2.6 ± 0.9	2.8 ± 0.6	0.71	2.5 ± 0.6	2.8 ± 0.9	0.16
HDL-C (mmol/l)	1.6 ± 0.4	1.4 ± 0.3	0.15	1.6 ± 0.3	1.4 ± 0.4	0.28	1.5 ± 0.4	1.6 ± 0.3	0.74
TG (mmol/l)	0.8 (0.6–0.9)	1.2 (0.9–1.7)	0.01	0.8 (0.6–1.1)	1.0 (0.7–1.2)	0.30	1.1 (0.8–1.6)	0.8 (0.6–0.9)	0.01
Lymphocytes (%)	38.4 ± 6.8	43.0 ± 8.5	0.08	42.2 ± 10.0	37.6 ± 4	0.09	42.1 ± 8.7	37.4 ± 8.4	0.11
Monocytes (%)	8.5 ± 1.8	8.1 ± 1.9	0.49	8.1 ± 1.7	8.6 ± 2.3	0.42	8.8 ± 2.2	8.48 ± 2.0	0.71
Plasma level of fatty (wt%)									
Total SFA	29.4 (28.8–30.5)	32.1 (31.1–33.2)	< 0.01	30.7 (29.3–32.3)	30.8 (29.8–31.8)	0.78	32.6 (31.5–33.2)	29.1 (28.7–29.5)	< 0.01
Myristic acid (14:0)	0.6 (0.6–1.0)	1.0 (0.8–1.2)	< 0.01	0.8 (0.6–1.1)	0.9 (0.7–1.1)	0.57	1.1 (0.9–1.2)	0.6 (0.6–0.9)	< 0.01
Palmitic acid (16:0)	20.5 (20.2–21.6)	23.4 (21.4–24.2)	< 0.01	20.9 (20.4–23.7)	21.5 (20.9–22.6)	0.39	23.9 (21.8–24.5)	20.4 (20.2–20.7)	< 0.01
Stearic acid (18:0)	8.1 ± 0.7	8.1 ± 1.1	0.86	8.2 ± 1.1	8.0 ± 1.0	0.65	8.0 ± 1.1	7.7 ± 0.8	0.37
Total n-6	38.6 ± 1.3	30.3 ± 2.9	< 0.01	31.0 ± 4.0	31.5 ± 4.5	0.74	30.8 ± 3.4	38.0 ± 2.0	< 0.01
LA (18:2n-6)	32.3 ± 1.6	24.9 ± 2.5	< 0.01	27.8 ± 3.8	29.2 ± 4.1	0.29	24.8 ± 2.5	31.8 ± 2.1	< 0.01
AA (20:4n-6)	6.2 ± 1.1	5.4 ± 0.9	0.02	5.8 ± 0.7	6.2 ± 1.2	0.32	5.9 ± 1.3	6.2 ± 1.1	0.58
Total n-3	5.8 ± 2.5	7.0 ± 2.3	0.12	9.0 ± 0.9	3.6 ± 0.49	< 0.01	6.1 ± 2.5	6.8 ± 2.4	0.44
ALA (18:3n-3)	0.5 (0.5–0.6)	0.5 (0.4–0.6)	0.46	0.5 (0.5–0.6)	0.5 (0.5–0.6)	0.79	0.5 (0.5–0.6)	0.5 (0.5–0.5)	0.95
EPA (20:5n-3)	1.7 (0.6–2.3)	2.4 (1.8–2.9)	0.14	3.2 (2.8–3.7)	0.6 (0.5–0.6)	< 0.01	2.0 (0.7–2.6)	2.2 (1.3–3.2)	0.46
DPA (22:5n-3)	0.6 ± 0.1	0.7 ± 0.2	0.26	0.8 ± 0.1	0.5 ± 0.1	< 0.01	0.6 ± 0.2	0.6 ± 0.1	0.80
DHA (22:6n-3)	3.0 (1.9–4.1)	4.0 (3.1–4.1)	0.14	4.3 (4.1–4.7)	1.9 (1.8–2.1)	< 0.01	3.5 (2.0–4.1)	3.3 (2.8–4.3)	0.48
Total PUFA	43.8 (42.1–47.3)	38.6 (35.8–39.7)	< 0.01	42.2 (39.5–46.7)	40.4 (37.7–41.5)	0.05	37.8 (35.8–39.1)	44.6 (43.1–47.3)	< 0.01
SFA to PUFA ratio	0.7 (0.6–0.7)	0.8 (0.8–0.9)	< 0.01	0.7 (0.6–0.8)	0.8 (0.7–0.8)	0.13	0.9 (0.8–0.9)	0.7 (0.6–0.7)	< 0.01

Lymphocytes and monocytes are given as percentage of total white cell count. Differences between tertiles were analyzed using the independent samples *t* test when normally distributed or the Mann-Whitney *U* test when not normally distributed. *P* values < 0.05 were considered significant. Values are presented as mean ± SD or medians and 25th–75th percentile

BMI body mass index, *Total-C* total cholesterol, *LDL-C* low-density lipoprotein cholesterol, *HDL-C* high-density lipoprotein cholesterol, *TG* triglycerides, *SFA* saturated fatty acid, *LA* linoleic acid, *AA* arachidonic acid, *ALA* alpha-linolenic acid, *EPA* eicosapentaenoic acid, *DPA* docosapentaenoic acid, *DHA* docosahexaenoic acid, *PUFA* polyunsaturated fatty acid